

ENHANCE RECOVERY WITH EVEN MORE THERAPEUTIC BENEFITS

DON'T JUST TREAT SYMPTOMS, HELP ACCELERATE HEALING

Upon recent Game Ready® specific post-market clinical follow-up studies, Game Ready is pleased to announce that there are nine additional benefits to claim, as well as further support of two existing performance claims (reduces swelling and patient's pain response). One of the most valuable clinical benefits is the reduction of patient narcotic pain medicine.

These claims further set Game Ready apart from competitors, including traditional RICE therapy, and demonstrate how patients can accelerate their recovery using the Game Ready System. Relieving pain and swelling is a start. But with Game Ready, you get the proactive power to help your patients heal faster and recover better.



GAME READY MODALITIES & BENEFITS

Game Ready's Active Compression with Active Cold Therapy

Reduces patient narcotic consumption and allows earlier discontinuation of narcotic pain medications.^{1,2,3,4}

Improves key, measurable physical therapy milestones.²

Increases patient satisfaction with recovery process.^{1,2}

Reduces likelihood for blood transfusion.¹

Shortens patient hospital admission time.^{1,5,6}

Provides less wound discharge.¹

Reduces risk of infection.¹

Improves surface contact for faster, deeper, longer-lasting cooling.^{5,6}

Decreases pain, muscle spasms, edema, and swelling.^{1,3,4,7,8}

Cold (General)

Reduces metabolic activity and cellular demand for oxygen, reducing secondary tissue damage.^{7,8}

Compression (General)

Aids the prevention of edema formation and removal of swelling.^{7,8,9}

Active Cold

Draws heat away from the injury site for more efficient temperature reduction.⁷

Active Compression

Mimics natural muscle contractions, "pumping" edema out of the injured area.^{7,8}

Optimizes lymphatic drainage.^{10,11}

Increases blood flow and oxygen delivery.^{12,13,14,15}

Stimulates tissue repair, encouraging acceleration of the healing process.^{7,8,12,13,14}

References:

1. Leegwater NC, Jore HW, Brohet R, Nolte P. Cryocompression therapy after elective arthroplasty of the hip. *Hip Int.* 2012;22(05):527-533.
2. Su EP, Perna M, Boettner F, Mayman DJ, Gerlinger T, Barsoum W, Randolph J, Lee G. A prospective, multi-center randomized trial to evaluate the efficacy of a cryopneumatic device on total knee arthroplasty recovery. *J Bone & Joint Surg.* November 2012. 94-B;SuppA:153-6.
3. Waterman W, Walker JJ, Swains C, Shortt M, Todd MD, Machen SM, Owens BD. (2012). The efficacy of combined cryotherapy and compression compared with cryotherapy alone following anterior cruciate ligament reconstruction. *J Knee Surg.* 2012;25(02):155-160.
4. Hochberg J. A randomized prospective study to assess the efficacy of two cold-therapy treatments following carpal tunnel release. *J Hand Surg.* 2001;July-Sept:208-215.
5. Womochel KS, Trowbridge CA, Keller D. The effects of continuous cooling and cyclical compression on intramuscular temperatures and cardiovascular strain. *Journal of Athletic Training.* 2009;44(3):s87.
6. Trowbridge CA, Womochel KS, and Davis DR. The effects of continuous cooling and cyclical compression on intramuscular and surface temperatures of the distal quadriceps. *Journal of Athletic Training.* 2008;43(3):S58.
7. Knight KL. Cryotherapy in sport injury management. 1995. *Human Kinetics, Champaign, IL.*
8. Sabiston KB et al. The effects of intermittent compression and cold on reducing edema in postacute ankle sprains. *Journal of Athletic Training.* 1992;27(2):140.
9. Capps/Mayberry. Cryotherapy and intermittent pneumatic compression for soft tissue trauma. January 2009.
10. McGeown JG, McHale NG, Thornbury KD. Effects of varying patterns of external compression on lymph flow in the hindlimb of the anaesthetized sheep. *J Physiol.* 1988;397:449-457.
11. McGeown JG, McHale NG, Thornbury KD. The role of external compression and movement in lymph propulsion in the sheep hind limb. *J Physiol.* 1987;387:83-93.
12. Hubbard TJ, Denegar CR. Does cryotherapy improve outcomes with soft tissue injury? *J Athl Train.* 2004;39(3):278-279.
13. Schaser K-D, Stover JF, Melcher I et al. Local cooling restores microcirculatory hemodynamics after closed soft-tissue trauma in rats. *J Trauma.* 2006;61(3):642-649.
14. Dahl J, Li J, Bring DK-I, Renström P, Ackerman PW. Intermittent pneumatic compression enhances neurovascular ingrowth and tissue proliferation during connective tissue healing. A study in the rat. *J Orthop Res.* 2007;25:1185-1192.
15. Johansson K, Lie E, Ekdahl C, Lindfeldt J. A randomized study comparing manual lymph drainage with sequential pneumatic compression for treatment of postoperative arm lymphedema. *Lymphology.* 1998;31(2):56-64.